


FPR Agonist 43

Chemical Properties

CAS No. :	903895-98-7
Formula:	C ₂₀ H ₂₁ ClN ₄ O ₂
Molecular Weight:	384.86
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	FPR Agonist 43 is a dual agonist of formyl peptide receptor 1 and formyl peptide receptor 2 (FPR2)/ALX.
Targets(IC50)	Others
In vitro	FPR Agonist 43 (10 ⁻⁵ -10 ⁻⁷ nM) is actively potent in the cAMP assay in FPR2/ALX over-expressing CHO cells. FPR Agonist 43 is also active in the GTPγ binding assay (IC ₅₀ =207±51 nM)[1]. FPR1 is the preferred receptor for FPR Agonist 43 in in both human neutrophils and possibly also in mouse cells[2].

Solubility Information

Solubility	DMSO: 34.2 mg/mL (88.86 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Corn Oil: 2 mg/mL (5.2 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.5983 mL	12.9917 mL	25.9835 mL
5 mM	0.5197 mL	2.5983 mL	5.1967 mL
10 mM	0.2598 mL	1.2992 mL	2.5983 mL
50 mM	0.052 mL	0.2598 mL	0.5197 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Planagumà A, et al. Lack of activity of 15-epi-lipoxin A₄ on FPR2/ALX and CysLT1 receptors in interleukin-8-driven human neutrophil function. Clin Exp Immunol. 2013 Aug;173(2):298-309.

Forsman H, et al. What formyl peptide receptors, if any, are triggered by compound 43 and lipoxin A₄? Scand J Immunol. 2011 Sep;74(3):227-234.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481