

## CP-809101 hydrochloride

## Chemical Properties


CAS No. : 1215721-40-6

Formula: C<sub>15</sub>H<sub>18</sub>Cl<sub>2</sub>N<sub>4</sub>O

Molecular Weight: 341.24

Storage: Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



## Biological Description

Description	CP-809101 hydrochloride is a potent and selective 5-HT <sub>2C</sub> receptor agonist with pEC <sub>50</sub> values of 9.96 for human 5-HT <sub>2C</sub> , 7.19 for 5-HT <sub>2B</sub> , and 6.81 for 5-HT <sub>2A</sub> receptors.
Targets(IC <sub>50</sub> )	5-HT Receptor
In vitro	CP-809101 is a potent, functionally selective 5-HT <sub>2C</sub> agonist that displays approximately 100% efficacy in vitro[4].
In vivo	Similar to currently available antipsychotic drugs, CP-809101 dose-dependently inhibited conditioned avoidance responding (CAR, ED <sub>50</sub> = 4.8 mg/kg, sc). CP-809101 antagonized both PCP- and d-amphetamine-induced hyperactivity with ED <sub>50</sub> values of 2.4 and 2.9 mg/kg (sc), respectively, and also reversed an apomorphine induced-deficit in prepulse inhibition. At doses up to 56 mg/kg, CP-809101 did not produce catalepsy. CP-809101 was inactive in two animal models of antidepressant-like activity, the forced swim test, and learned helplessness[4].

## Solubility Information

Solubility	H <sub>2</sub> O: 20 mg/mL (58.61 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--

### Preparing Stock Solutions

---

	1mg	5mg	10mg
1 mM	2.9305 mL	14.6524 mL	29.3049 mL
5 mM	0.5861 mL	2.9305 mL	5.861 mL
10 mM	0.293 mL	1.4652 mL	2.9305 mL
50 mM	0.0586 mL	0.293 mL	0.5861 mL

---

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Higgins GA, Silenieks LB, Lau W, et al. Evaluation of chemically diverse 5-HT<sub>2C</sub> receptor agonists on behaviours motivated by food and nicotine and on side effect profiles. *Psychopharmacology (Berl)*. 2013 Apr;226(3):475-90.
- Strong PV, Christianson JP, Loughridge AB, et al. 5-hydroxytryptamine 2C receptors in the dorsal striatum mediate stress-induced interference with negatively reinforced instrumental escape behavior. *Neuroscience*. 2011 Dec 1; 197:132-44. doi: 10.1016/j.neuroscience.2011.09.041. Epub 2011 Sep 24.
- Fletcher PJ, Tampakeras M, Sinyard J et al. Characterizing the effects of 5-HT(2C) receptor ligands on motor activity and feeding behaviour in 5-HT(2C) receptor knockout mice. *Neuropharmacology*. 2009 Sep;57(3):259-67. doi: 10.1016/j.neuropharm.2009.05.011.
- Siuciak JA, Chapin DS, McCarthy SA, et al. CP-809,101, a selective 5-HT<sub>2C</sub> agonist, shows activity in animal models of antipsychotic activity. *Neuropharmacology*. 2007 Feb;52(2):279-90.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481