

Ac-DEVD-CHO

Chemical Properties


CAS No. : 169332-60-9

Formula: C₂₀H₃₀N₄O₁₁

Molecular Weight: 502.47

Storage: Keep away from moisture, Store at low temperature
Powder: -20°C for 3 years | In solvent: -80°C for 1 year

Actual storage temperature shall be subject to the COA.



Biological Description

Description	Ac-DEVD-CHO is a specific Caspase-3 inhibitor (K _i : 230 pM) with inhibitory effects on SLNT-induced apoptosis.
Targets(IC ₅₀)	Apoptosis, Caspase
In vitro	The addition of Ac-DEVD-CHO significantly prevents SLNT-induced apoptosis (from 32.91% decreases to 15.88% while NC and Ac-DEVD-CHO groups are 6.45%, and 7.77%, respectively) [2]. Ac-DEVD-CHO (10 μmol/L) partially blocks the effect of SIN-induced apoptosis and reduces the number of apoptotic nuclei. These effects of SIN are blocked by the caspase-3 inhibitor Ac-DEVD-CHO. Camptothecin (4 μM), a positive control, increases caspase-3 activity, which is also blocked by Ac-DEVD-CHO.[3]
In vivo	One hundred and two male C57BL/6 mice were subjected to cecal ligation and puncture (CLP, a model of polymicrobial sepsis) or sham operation. The animals were assigned into three equal groups (n=34) according to the random number table: sham group, model group, and caspase-3 inhibitor (CI) group. Thirty minutes before CLP, Ac-DEVD-CHO (4 μg/g) was injected subcutaneously in the CI group. The levels of blood urea nitrogen (BUN) and creatinine (Cr) were determined, and the concentrations of tumor necrosis factor-α (TNF-α), interleukins (IL-6 and IL-10) were measured by enzyme-linked immunosorbent assay (ELISA), the renal cell apoptosis rate was determined by flow cytometry and the expression of caspase-3 mRNA was determined by real-time reverse transcription-polymerase chain reaction (RT-PCR) at 6, 12 and 24 hours after operation. The 4-day and 7-day survival rates of three groups of mice were observed. Results: Compared with the sham group, the concentrations of serum BUN, TNF-α, IL-6, IL-10, and the renal cell apoptosis rates, the caspase-3 mRNA expression were increased significantly at all time points after CLP, the concentrations of serum Cr were increased significantly at 6 hours, with the 4-day and 7-day survival rates were decreased significantly. Compared with the model group, in the CI group, the concentrations of serum BUN were decreased significantly at all time points after operation and those of Cr were decreased significantly at 6 hours, then restored to those of the sham group at 12 hours and 24 hours; the concentrations of serum TNF-α, IL-6 were decreased and those of IL-10 elevated significantly at all time points. (P<0.05); the renal cell apoptosis rate and the expression of caspase-3 mRNA were decreased significantly at all time points. The 4-day survival rate of the CI group was improved (80% vs. 20%), but that of

In vivo	the 7-day did not change (20% vs. 20%).[4]
---------	--------------------------------------------

Solubility Information

Solubility	H2O: 255 mg/mL (507.49 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
------------	--------------------------------------------------------------------------------------------------------------------------

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.9902 mL	9.9508 mL	19.9017 mL
5 mM	0.398 mL	1.9902 mL	3.9803 mL
10 mM	0.199 mL	0.9951 mL	1.9902 mL
50 mM	0.0398 mL	0.199 mL	0.398 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Garcia-Calvo M, et al. Inhibition of human caspases by peptide-based and macromolecular inhibitors. *J Biol Chem.* 1998 Dec 4;273(49):32608-13.

Jinglin Wang, et al. A polysaccharide from *Lentinus edodes* inhibits human colon cancer cell proliferation and suppresses tumor growth in athymic nude mice. *Oncotarget.* 2017 Jan 3; 8(1): 610-623.

Long-gang He, et al. Sinomenine induces apoptosis in RAW 264.7 cell-derived osteoclasts in vitro via caspase-3 activation. *Acta Pharmacol Sin.* 2014 Feb; 35(2): 203-210.

Liu LX, et al. The effect of caspase-3 inhibitor on the concentrations of serum inflammatory cytokines in sepsis related acute kidney injury induced by peritoneal cavity infection in mice. *Zhongguo Wei Zhong Bing Ji Jiu Yi Xue.* 2010 Dec;22(12):736-9.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481