


## Paclitaxel

### Chemical Properties

CAS No. :	33069-62-4
Formula:	C <sub>47</sub> H <sub>51</sub> NO <sub>14</sub>
Molecular Weight:	853.91
Storage:	Keep away from direct sunlight, Store at low temperature Powder: -20°C for 3 years   In solvent: -80°C for 1 year <small>Actual storage temperature shall be subject to the COA.</small>



### Biological Description

Description	Paclitaxel (Taxol) is a natural product and a microtubule polymer stabilizer. Paclitaxel has anti-tumor activity and causes cell death by inducing mitotic arrest, apoptosis, and cell autophagy.
Targets(IC50)	Apoptosis, Microtubule Associated, Autophagy, ADC Cytotoxin
In vitro	<p><b>METHODS:</b> Human T-cell lymphoblastic leukemia cells CCRF-HSB-2 were treated with Paclitaxel (0.01-1 μM) for 48 h. Cell growth inhibition was detected by MTT.</p> <p><b>RESULTS:</b> Paclitaxel dose-dependently inhibited the growth of CCRF-HSB-2 cells with an IC<sub>50</sub> of 0.25 μM. [1]</p> <p><b>METHODS:</b> Human gastric cancer cells AGS were treated with Paclitaxel (10-160 nM) for 24-48 h. The expression levels of target proteins were detected by Western Blot.</p> <p><b>RESULTS:</b> Paclitaxel induced the up-regulation of the expression of cleaved caspase-3 and cleaved PARP, which are apoptosis-related proteins. [2]</p> <p><b>METHODS:</b> Canine mammary tumor cells CHMm were treated with Paclitaxel (0.01-1 μM) for 24 h. Apoptosis was detected using Flow Cytometry.</p> <p><b>RESULTS:</b> Paclitaxel dose-dependently induced apoptosis in CHMm cells. [3]</p>
In vivo	<p><b>METHODS:</b> To investigate the effect of low-dose Paclitaxel on tumor invasion, Paclitaxel (2.6 mg/kg) was administered intraperitoneally to SCID mice bearing cholangiocarcinoma tumor EGI-1 once daily for two weeks.</p> <p><b>RESULTS:</b> Low-dose Paclitaxel treatment reduced the pulmonary spread of EGI-1 cells without significantly affecting their local tumor growth. [4]</p> <p><b>METHODS:</b> To develop a preclinical model of Paclitaxel alcohol-induced negative affective symptoms, Paclitaxel (2-8 mg/kg in 1 volume ethanol+1 volume Emulphor-620 +18 volumes distilled water) was injected intraperitoneally into C57BL/6J mice every four injections were given to C57BL/6J mice once every two days.</p> <p><b>RESULTS:</b> 8 mg/kg Paclitaxel treatment resulted in the development and maintenance of mechanical and cold abnormalities of pain. Paclitaxel also induced anxiety-like and depressive-like behaviors. Paclitaxel produced behavioral changes in the mouse affective state modeling assay, whereas the increase in injurious responses lasted longer. [5]</p>
Kinase Assay	To determine which caspases are involved in apoptosis induced by taxol, caspase-3 inhibitor (DEVD-CHO), caspase-6 inhibitor (Z-VEID-FMK), caspase-8 inhibitor (Z-IETD-

Kinase Assay	FMK or IETD-CHO), caspase-9 inhibitors (Z-LEHD-FMK or LEHD-CHO), and caspase-10 inhibitor (Z-AEVD-FMK) are used. These caspase inhibitors are dissolved in dimethyl sulfoxide (Me2SO); the final concentration of Me2SO is 0.1%. Cells (5×10 <sup>5</sup> ) are preincubated in the presence or absence of 100 μM each of these inhibitors for 3 h at 37°C then treated with or without 0.1, 0.5, and 1 μM Paclitaxel for 48 h and processed for annexin V binding assay [1].
Cell Research	1×10 <sup>4</sup> cells are plated in 100 μL of the growth medium in the presence or absence of increasing concentrations (0.1–1 μM) of taxol in 96-well plates and cultured at 37°C in 5% CO <sub>2</sub> for 12–48 h. The cells are then incubated with 25 μL of MTT (5 mg/mL) at 37°C for 4 h. After dissolving the crystals with 0.04 N HCl in isopropanol, the plates are read in a microplate reader at 570 nm [1].
Animal Research	Adult (250–320 g) male Sprague-Dawley rats are used for all experiments. One week following the Dil injection, rats are anesthetized with isoflurane and injected into the tail vein with 2 mg/kg paclitaxel or its vehicle (1:1:23, cremophor EL:ethanol:0.9% saline). The tail vein injection is repeated three more times every other day for a total of four injections [4].

### Solubility Information

Solubility	Ethanol: 21.4 mg/mL (25.06 mM), Sonication is recommended. DMSO: 257.5 mg/mL (301.55 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 8.54 mg/mL (10 mM), Suspension. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

### Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	1.1711 mL	5.8554 mL	11.7108 mL
5 mM	0.2342 mL	1.1711 mL	2.3422 mL
10 mM	0.1171 mL	0.5855 mL	1.1711 mL
50 mM	0.0234 mL	0.1171 mL	0.2342 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

### Reference

- Park SJ, et al. Taxol induces caspase-10-dependent apoptosis. *J Biol Chem*. 2004 Dec 3;279(49):51057-67.
- Gritsina G, Fong K, Lu X, et al. Chemokine receptor CXCR7 activates AURKA and promotes neuroendocrine prostate cancer growth. *The Journal of Clinical Investigation*. 2023
- Zeng L, Lyu X, Yuan J, et al. STMN1 Promotes Tumor Metastasis in Non-small Cell Lung Cancer Through Microtubule-dependent And Nonmicrotubule-dependent Pathways. *International Journal of Biological Sciences*. 2024, 20(4): 1509.
- Lin Y, Chen X, Lin L, et al. Sesamolin serves as an MYH14 inhibitor to sensitize endometrial cancer to chemotherapy and endocrine therapy via suppressing MYH9/GSK3 $\beta$ / $\beta$ -catenin signaling. *Cellular & Molecular Biology Letters*. 2024, 29(1): 63.
- Bao Q, Wang A, Hong W, et al. The c-Abl-RACK1-FAK signaling axis promotes renal fibrosis in mice through regulating fibroblast-myofibroblast transition. *Cell Communication and Signaling*. 2024, 22(1): 247.
- Gao X, Ren X, Wang F, et al. Immunotherapy and drug sensitivity predictive roles of a novel prognostic model in hepatocellular carcinoma. *Scientific Reports*. 2024, 14(1): 9509.
- Gonzalez-Posada A H, Mesa M, Sierra L, et al. Interactions of human serum albumin with phosphate and Tris buffers: impact on paclitaxel binding and nanoparticles self-assembly. *Journal of Microencapsulation*. 2024: 1-12.
- Meng P, Liu W, Lao J, et al. Paclitaxel improves thrombopoiesis in the absence of thrombopoietin receptor (Mpl). *Journal of Thrombosis and Haemostasis*. 2024
- Wang J, Liu P, Zhang R, et al. VASH2 enhances KIF3C-mediated EGFR-endosomal recycling to promote aggression and chemoresistance of lung squamous cell carcinoma by increasing tubulin dephosphorylation. *Cell Death & Disease*. 2024, 15(10): 772.
- Sun L Y, Ke S B, Li B X, et al. ANP32E promotes esophageal cancer progression and paclitaxel resistance via P53/SLC7A11 axis-regulated ferroptosis. *International Immunopharmacology*. 2025, 144: 113436.
- Guo K, Gao L, Li P, et al. Allopregnanolone relieves paclitaxel induced mechanical hypersensitivity via inhibiting spinal cord PGE2-EP2 mediated microglia-neuron signaling. *IBRO Neuroscience Reports*. 2025
- Zhao X, Lian X, Xie J, et al. Accumulated cholesterol protects tumours from elevated lipid peroxidation in the microenvironment. *Redox Biology*. 2023: 102678.
- Khing TM, et al. The effect of paclitaxel on apoptosis, autophagy and mitotic catastrophe in AGS cells. *Sci Rep*. 2021 Dec 6;11(1):23490.
- Ren X, et al. Paclitaxel suppresses proliferation and induces apoptosis through regulation of ROS and the AKT/MAPK signaling pathway in canine mammary gland tumor cells. *Mol Med Rep*. 2018 Jun;17(6):8289-8299.
- Zhu X, Huang N, Ji Y, et al. Brusatol induces ferroptosis in oesophageal squamous cell carcinoma by repressing GSH synthesis and increasing the labile iron pool via inhibition of the NRF2 pathway. *Biomedicine & Pharmacotherapy*. 2023, 167: 115567.
- Hou Z, Lin S, Du T, et al. S-72, a Novel Orally Available Tubulin Inhibitor, Overcomes Paclitaxel Resistance via Inactivation of the STING Pathway in Breast Cancer. *Pharmaceuticals*. 2023, 16(5): 749.
- Cadamuro M, et al. Low-Dose Paclitaxel Reduces S100A4 Nuclear Import to Inhibit Invasion and Hematogenous Metastasis of Cholangiocarcinoma. *Cancer Res*. 2016 Aug 15;76(16):4775-84.
- Zhan Y, Chen Q, Song Y, et al. Berbamine Hydrochloride inhibits lysosomal acidification by activating Nox2 to potentiate chemotherapy-induced apoptosis via the ROS-MAPK pathway in human lung carcinoma cells. *Cell Biology and Toxicology*. 2022: 1-21
- Toma W, et al. Effects of paclitaxel on the development of neuropathy and affective behaviors in the mouse. *Neuropharmacology*. 2017 May 1;117:305-315.
- Li Q, et al. Low doses of paclitaxel enhance liver metastasis of breast cancer cells in the mouse model. *FEBS J*. 2016 Aug;283(15):2836-52.
- Pang G, Zhang S, Zhou X, et al. Immunoactive polysaccharide functionalized gold nanocomposites promote dendritic cell stimulation and antitumor effects. *Nanomedicine*. 2019 May;14(10):1291-1306
- Sun J, Liu W, Li Y, et al. An on-chip cell culturing and combinatorial drug screening system[J]. *Microfluidics and Nanofluidics*. 2017 Jul 21(7): 125.
- Sun J, Liu W, Li Y, et al. An on-chip cell culturing and combinatorial drug screening system. *Microfluidics and Nanofluidics*. 2017 Jul 21(7): 125.
- Jing C, et al. Lenvatinib enhances the antitumor effects of paclitaxel in anaplastic thyroid cancer. *Am J Cancer Res*.

2017 Apr 1;7(4):903-912.

ZeZhanga,b,c VLM catecholaminergic neurons control tumor growth by regulating CD8+ T cells. PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA. 2020

Song X, Li D, Gan L, et al. Intravenous Injection of Na Ions Aggravates Ang II-Induced Hypertension-Related Vascular Endothelial Injury by Increasing Transmembrane Osmotic Pressure. International Journal of Nanomedicine. 2023: 7505-7521.

Pang G, Zhang S, Zhou X, et al. Immunoactive polysaccharide functionalized gold nanocomposites promote dendritic cell stimulation and antitumor effects[J]. Nanomedicine. 2019 May;14(10):1291-1306.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481