


Wnt pathway activator 2

Chemical Properties

CAS No. :	1360540-82-4
Formula:	C17H15NO4
Molecular Weight:	297.31
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Wnt pathway activator 2, with an IC50 of 13 nM, is a potent Wnt activator.
Targets(IC50)	Wnt/beta-catenin

Solubility Information

Solubility	DMSO: 90 mg/mL (302.71 mM), Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+90% Saline: 5 mg/mL (16.82 mM), Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	3.3635 mL	16.8175 mL	33.6349 mL
5 mM	0.6727 mL	3.3635 mL	6.727 mL
10 mM	0.3363 mL	1.6817 mL	3.3635 mL
50 mM	0.0673 mL	0.3363 mL	0.6727 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Charlene F. Barroga, et al. Diketones and hydroxyketones as catenin signaling pathway activators. WO2012024404A1.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481