


## Artesunate

## Chemical Properties

CAS No. :	88495-63-0
Formula:	C <sub>19</sub> H <sub>28</sub> O <sub>8</sub>
Molecular Weight:	384.42
Storage:	Powder: -20°C for 3 years   In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



## Biological Description

Description	Artesunate (WR-256283) is part of the artemisinin group of drugs that treat malaria. It is a semi-synthetic derivative of artemisinin, Artesunate is soluble in water but has poor stability in aqueous solutions at neutral or acid pH.
Targets(IC50)	Ferroptosis,STAT,Parasite,Virus Protease
In vitro	ART could alter the biomechanical properties of the glioma cells to inhibit cell proliferation, migration and invasion [2]. The ART significantly suppresses the cell proliferation, induces the apoptosis and promoted cell cycle arrest at G <sub>0</sub> /G <sub>1</sub> phase in SKM-1 cells. The mechanisms of ART anti-MDS is associated with the increase of intracellular calciumion concentration and ROS levels. In addition, the pro-apoptotic activity of ART may be involved in the regulation of BCL-2 /BAX ratio and the expressions of P-bad and survivin[3]. ART treatment demethylates CDH1, which, in turn recovers the E-cadherin activation in SKM-1 cells[1].
In vivo	Artesunate is a medication used to treat malaria. Artesunate can induce radiosensitivity of HeLa cells in vivo using Xenograft model of nude mice[5]. Artesunate had a relatively high immunosuppressive activity with low toxicity, and could inhibit T lymphocyte proliferation induced by mitogen and alloantigen[6].
Cell Research	Growth inhibition assay: The SKM-1 cells (1×10 <sup>5</sup> /mL) are firstly seeded in 96-well plates. Artesunate is diluted in 0.1% dimethyl sulfoxide (DMSO) producing 0, 12.5, 25, 50µg/mL concentrations and added to the SKM-1 cells with 100 µl per well. A negative control is treated with 0.1% DMSO. At 0, 24, 48, and 72 hours, same amount of MTT solution is added into each well and cultured for extra 4 hours. MTT treated cells are fixed with 150 µL DMSO for 30 min at room temperature and then determined with Evolution 201 and 220 UV-Vis spectrophotometer system at 540 nm. (Only for Reference)

## Solubility Information

Solubility	Ethanol: 9 mg/mL (23.41 mM),Sonication is recommended. H <sub>2</sub> O: < 1 mg/mL (insoluble or slightly soluble), DMSO: 250 mg/mL (650.33 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 5 mg/mL (13.01 mM),Sonication is recommended.

In vivo Formulation	<i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one. Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>
---------------------	--

**Preparing Stock Solutions**

	<b>1mg</b>	<b>5mg</b>	<b>10mg</b>
1 mM	2.6013 mL	13.0066 mL	26.0132 mL
5 mM	0.5203 mL	2.6013 mL	5.2026 mL
10 mM	0.2601 mL	1.3007 mL	2.6013 mL
50 mM	0.052 mL	0.2601 mL	0.5203 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

**Reference**

Xu N, et al. Int J Med Sci. 2015, 12(6):524-529.

Liu Y, Li H, Luo Z, et al. Artesunate, a new antimalarial clinical drug, exhibits potent anti-AML activity by targeting the ROS/Bim and TFRC/Fe2+ pathways. British Journal of Pharmacology. 2022

Jin J, Guo D, Wang Y, et al. Artesunate Inhibits the Development of Esophageal Cancer by Targeting HK1 to Reduce Glycolysis Levels in Areas With Zinc Deficiency. Frontiers in Oncology. 2022: 1956

Lian S, et al. Oncol Rep. 2016, 36(2):984-1990.

Qiao SK, et al. Zhongguo Shi Yan Xue Ye Xue Za Zhi. 2016, 24(1):131-137.

Bowden G D, Land K M, O'Connor R M, et al. High-throughput screen of drug repurposing library identifies inhibitors of Sarcocystis neurona growth. International Journal for Parasitology: Drugs and Drug Resistance. 2018 Apr; 8(1): 137-144

Wang J, Zhang J, Guo Z, et al. Targeting HSP70 chaperones by rhein sensitizes liver cancer to artemisinin derivatives. Phytomedicine. 2023: 155156.

Lisewski AM, et al. Cell. 2014, 158(4):916-928.

Luo J, et al. Radiat Oncol. 2014, 9:84.

Han X, Jiang S, Hu C, et al. Inhibition of Keloid Fibroblast Proliferation by Artesunate is Mediated by Targeting the IRE1 $\alpha$ /XBP1 Signaling Pathway and Decreasing TGF- $\beta$ 1. Burns. 2024

Wang B, et al. Artesunate sensitizes ovarian cancer cells to cisplatin by downregulating RAD51. Cancer Biol Ther. 2015;16(10):1548-56.

Ilamathi M, et al. Artesunate as an Anti-Cancer Agent Targets Stat-3 and Favorably Suppresses Hepatocellular Carcinoma. Curr Top Med Chem. 2016;16(22):2453-63.

Bowden G D, Land K M, O'Connor R M, et al. High-throughput screen of drug repurposing library identifies inhibitors of Sarcocystis neurona growth[J]. International Journal for Parasitology: Drugs and Drug Resistance. 2018 Apr; 8(1): 137-144.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

**This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use**

Tel: 781-999-4286 E\_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481