


Primaquine diphosphate

Chemical Properties

CAS No. :	63-45-6
Formula:	C ₁₅ H ₂₁ N ₃ O-2H ₃ PO ₄
Molecular Weight:	455.34
Storage:	Powder: -20°C for 3 years In solvent: -80°C for 1 year Actual storage temperature shall be subject to the COA.



Biological Description

Description	Primaquine diphosphate is the phosphate salt form of primaquine, a synthetic, 8-aminoquinoline derivative with antimalarial properties. Although its mechanism of action is unclear, primaquine bind to and alter the properties of protozoal DNA. This agent eliminates tissue (exo-erythrocytic) malarial infection, preventing the development of the erythrocytic forms of the parasite which are responsible for relapses in Plasmodium vivax and ovale malaria. Primaquine is active against late hepatic stages (hypnozoites, schizonts).
Targets(IC50)	Antibiotic,Parasite
In vivo	Primaquine effectively counters all exoerythrocytic forms of parasites and is used in conjunction with other antimalarial drugs to treat vivax and ovale malaria. It remains an efficacious malaria-blocking agent, exhibiting notable activity against the gametocytes of all human malaria, including multidrug-resistant strains of P. falciparum. The specific mechanism by which Primaquine eliminates P. falciparum and gametocytes is unclear, but studies suggest it involves disruptions to the parasite's mitochondrial metabolism, impairing coenzyme Q's role as an electron carrier in the respiratory chain, and potential generation of highly reactive metabolites leading to increased oxidative potential. Primaquine can block calcium-release-activated channels in rat megakaryocytes and may also act as an inhibitor of vesicular transport. It inhibits protein transport (IC ₅₀ : 50 μM) and vesicle budding in donor cell membranes. Primaquine significantly suppresses the recycling of phagocytosed proteins to the plasma membrane and inhibits potassium channel currents by blocking muscarinic acetylcholine receptors. It also inhibits human erythrocyte membrane acetylcholinesterase (IC ₃₃ : 30 μM) and affects the sodium channels in isolated rat

Solubility Information

Solubility	DMSO: 245 mg/mL (538.06 mM),Sonication is recommended. Ethanol: < 1 mg/mL (insoluble or slightly soluble), H ₂ O: 83 mg/mL (182.28 mM),Sonication is recommended. (< 1 mg/ml refers to the product slightly soluble or insoluble)
In vivo Formulation	10% DMSO+40% PEG300+5% Tween 80+45% Saline: 2 mg/mL (4.39 mM),Sonication is recommended. <i>Please add the solvents sequentially, clarifying the solution as much as possible before adding the next one.</i>

A DRUG SCREENING EXPERT

In vivo Formulation	<i>Dissolve by heating and/or sonication if necessary. Working solution is recommended to be prepared and used immediately. The formulation provided above is for reference purposes only. In vivo formulations may vary and should be modified based on specific experimental conditions.</i>
---------------------	--

Preparing Stock Solutions

	1mg	5mg	10mg
1 mM	2.1962 mL	10.9808 mL	21.9616 mL
5 mM	0.4392 mL	2.1962 mL	4.3923 mL
10 mM	0.2196 mL	1.0981 mL	2.1962 mL
50 mM	0.0439 mL	0.2196 mL	0.4392 mL

Please select the appropriate solvent to prepare the stock solution, according to the solubility of the product in different solvents. Please use it as soon as possible.

Note: The dilution table applies only to solid products. For liquid products, please calculate the stock solution based on the stated concentration and/or density.

Reference

Vale N, et al. Eur J Med Chem, 2009, 44(3), 937-953.

Bowden G D, Land K M, O'Connor R M, et al. High-throughput screen of drug repurposing library identifies inhibitors of Sarcocystis neurona growth. International Journal for Parasitology: Drugs and Drug Resistance. 2018 Apr; 8(1): 137-144

Ding L, Chen X, Zhang W, et al. Canagliflozin primes antitumor immunity by triggering PD-L1 degradation in endocytic recycling. The Journal of Clinical Investigation. 2023, 133(1).

Bowden G D, Land K M, O'Connor R M, et al. High-throughput screen of drug repurposing library identifies inhibitors of Sarcocystis neurona growth. International Journal for Parasitology: Drugs and Drug Resistance. 2018 Apr; 8(1): 137-144.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481