

GABARAP Protein, Human, Recombinant (His & hFc)

General Information

Synonyms:	GABA(A) Receptor-Associated Protein;HCG1987397 Isoform CRA_b;GABARAP Protein;GABARAP
Protein Construction:	Met1-Leu117
Species:	Human
Expression Host:	E. coli
Accession:	Q6IAW1
Molecular Weight:	50 KDa (reducing condition)
AA Sequence:	Met1-Leu117

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Supplied as a 0.2 μm filtered solution of 20 mM PB, 150 mM NaCl, 20% Glycerol, pH 7.0.

Preparation and Storage

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

Proteins are shipped with blue ice.

Protein Background

Gamma-Aminobutyric Acid Receptor-Associated Protein (GABARAP) is a ligand-gated chloride channel protein that mediates inhibitory neurotransmission and belongs to the MAP1 LC3 family. GABARAP is highly positively charged in its N-terminus and shares sequence similarity with light chain-3 of microtubule-associated proteins 1A and 1B. GABARAP clusters neurotransmitter receptors by mediating interaction with the cytoskeleton. Autophagy is the process by which cells recycle cytoplasm and dispose of excess or defective organelles. This process is suggested to be involved development, differentiation, growth regulation and tissue remodeling in multicellular organisms. An important inhibitory neurotransmitter, GABA, acts on GABA receptors that are ubiquitously expressed in the CNS. GABARAP has been shown to play a important role in intracellular transport of GABA(A) receptors and its interaction with the cytoskeleton.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481