

IL-11 Protein, Human, Recombinant (P. pastoris)

General Information

Synonyms:	Oprelvekin;Adipogenesis Inhibitory Factor;IL11;AGIF;IL-11;Interleukin-11
Protein Construction:	Gly23-Leu199
Species:	Human
Expression Host:	P. pastoris (Yeast)
Accession:	P20809
Molecular Weight:	19 KDa (reducing condition)
AA Sequence:	Gly23-Leu199

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 2% Glycine, pH 7.2.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Interleukin 11 (IL-11) is a member of a family of human growth factors that includes human growth hormone, granulocyte colony-stimulating factor, and other growth factors. IL-11 is a thrombopoietic growth factor that directly stimulates the proliferation of hematopoietic stem cells and megakaryocyte progenitor cells and induces megakaryocyte maturation resulting in increased platelet production. It also promotes the proliferation of hepatocytes in response to liver damage. Binding to its receptor formed by IL6ST and either IL11RA1 or IL11RA2, It

A DRUG SCREENING EXPERT

activates a signaling cascade that promotes cell proliferation. The signaling leads to the activation of intracellular protein kinases and the phosphorylation of STAT3.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481