

4-1BB/CD137/TNFRSF9 Protein, Human, Recombinant (hFc & His)

General Information

Synonyms:	CD137;4-1BB ligand receptor;T-cell antigen 4-1BB homolog;T-cell antigen ILA;ILA;CDw137;TNFRSF9
Protein Construction:	Leu24-Gln186
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q07011
Molecular Weight:	55-75 KDa (reducing condition)
AA Sequence:	Leu24-Gln186

QC Testing

Biological Activity:	1. Loaded Human 4-1BB-Fc-His on HIS1K Biosensor, can bind Anti-Human CD137 mAb with an affinity constant of 12.2 nM as determined in BLI assay. (Regularly tested) 2. Immobilized Human 4-1BB-Fc-His at 2µg/ml (100 µl/well) can bind Anti-Human CD137 mAb. The ED50 of Anti-Human CD137 mAb is 0.41 µg/ml. (Regularly tested)
Endotoxin:	< 0.1 ng/µg (1 EU/µg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Tumor necrosis factor receptor superfamily member 9 (TNFRSF9) is an inducible T cell surface protein belonging to the TNF receptor superfamily. It is a single-pass type I membrane protein which contains 4 TNFR-Cys repeats. The human and mouse proteins share 60% amino acid sequence identity. It is absent from naive T cells, but upregulated and continually expressed following T cell activation. It is a receptor for TNFSF9/4-1BBL, and possibly

active during T cell activation.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481