

IL-20RB Protein, Human, Recombinant (hFc)

General Information

| | |
|-----------------------|---|
| Synonyms: | DIRS1;IL-20R- β ;Interleukin-20 receptor subunit β ;IL-20 receptor subunit beta;IL-20RB;interleukin-20 receptor II;hCG_2022374;fibronectin type III domain containing 6;IL-20 receptor subunit β ;Interleukin-20 receptor subunit beta;MGC34923;FNDC6;IL-20R-beta;IL-20R2 |
| Protein Construction: | Asp30-Ala230 |
| Species: | Human |
| Expression Host: | HEK293 Cells |
| Accession: | Q6UXL0 |
| Molecular Weight: | 60-85 KDa (reducing condition) |
| AA Sequence: | Asp30-Ala230 |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | Greater than 90% as determined by reducing SDS-PAGE. (QC verified) |
| Endotoxin: | < 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test. |
| Formulation: | Supplied as a 0.2 μ m filtered solution of 20 mM PB, 150 mM NaCl, pH 7.4. |

Preparation and Storage

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

Proteins are shipped with blue ice.

Protein Background

Interleukin-20 receptor subunit beta (IL20RB) is a single-pass type I membrane protein and belongs to the type II cytokine receptor family. It contains 2 fibronectin type-III domains. There are two kinds of type II cytokine receptors : cytokine receptors that bind type I and type II interferons; cytokine receptors that bind members of the interleukin-10 family (interleukin-10, interleukin-20 and interleukin-22). Type II cytokine receptors are similar to type I cytokine receptors except they do not possess the signature sequence WSXWS that is characteristic of type I receptors. They are expressed on the surface of certain cells, which bind and respond to a select group of cytokines. These receptors are related predominantly by sequence similarities in their extracellular portions that

A DRUG SCREENING EXPERT

are composed of tandem Ig-like domains. The intracellular domain of type II cytokine receptors is typically associated with a tyrosine kinase belonging to the Janus kinase (JAK) family.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481