

Fas/CD95 Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	CD95;Apoptosis-mediating surface antigen FAS;TNFRSF6;Fas;Tumor necrosis factor receptor superfamily member 6;FASLG receptor;Apo-1 antigen
Protein Construction:	Gln22-Arg169
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P25446
Molecular Weight:	55 KDa (reducing condition)
AA Sequence:	Gln22-Arg169

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Mouse Apoptosis-mediating surface antigen FAS (Fas) belongs to the death receptor subfamily of the TNF receptor superfamily and is designated TNFRSF6. Mouse Fas contains 1 death domain and 3 TNFR-Cys repeats. It detected in various tissues including thymus, liver, lung, heart, and adult ovary. As a receptor for TNFSF6/FASLG, The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases

mediating apoptosis. FAS-mediated apoptosis may have a role in the induction of peripheral tolerance, in the antigen-stimulated suicide of mature T-cells, or both.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481