

SLAMF1 Protein, Mouse, Recombinant (His)

General Information

| | |
|-----------------------|--|
| Synonyms: | signaling lymphocytic activation molecule;SLAM family member 1;CD150 antigen;SLAMF1;SLAM;CD150 |
| Protein Construction: | Thr25-Pro242 |
| Species: | Mouse |
| Expression Host: | HEK293 Cells |
| Accession: | Q9QUM4 |
| Molecular Weight: | 40-60 KDa (reducing condition) |
| AA Sequence: | Thr25-Pro242 |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first. |
| Purity: | Greater than 95% as determined by reducing SDS-PAGE. (QC verified) |
| Endotoxin: | < 0.1 ng/μg (1 EU/μg) as determined by LAL test. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4. |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Signaling lymphocyte activation molecule (SLAM), is a self-ligand glycoprotein which exists not only found on the surface of activated and memory T cells, but also on the surface of activated B cells, dendritic cells, and macrophages. SLAM consists of a extracellular domain (ECD) with two Ig-like domains,transmembrane segment, and cytoplasmic domain with three immunoreceptor tyrosine switch motifs (ITSM). SLAM is thought to play an important role in adhesion between T cells and APCs and has been shown to act as a coreceptor in TCR-dependent

responses. SLAM, together with CD46, is one of the two receptors for measles virus. SLAM is a cell surface receptor that, like the B cell receptor, CD40, and CD95, can transmit positive or negative signals. SLAM can associate with the SH2-containing inositol phosphatase (SHIP), the SH2-containing protein tyrosine phosphatase (SHP-2), and the adaptor protein SH2 domain protein 1A. It's upregulated on activated B cells and CD4+ and CD8+ T cells, but downregulated on Th2 polarized cells. Also, it can Inhibits antigen receptor-mediated production of IFN-gamma, but not IL-2, in CD4-/CD8- T-cells.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481