

TRAIL R2/DR5/TNFRSF10B Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	MK;CD262;Tnfrsf10b;Dr5;Tumor necrosis factor receptor superfamily member 10B;Killer; Death receptor 5
Protein Construction:	Asn53-Ser177
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q9QZM4
Molecular Weight:	50-75 KDa (reducing condition)
AA Sequence:	Asn53-Ser177

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 150 mM NaCl, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Mouse tumor necrosis factor receptor superfamily member 10B (TNFRSF10B) is a member of the TNFR family which contains 1 death domain and 3 TNFR-Cys repeats. TNFRSF10B exhibits high structural and functional homology to TRAIL-R1 (DR-4). TNFRSF10B is highly expressed in heart, lung, lymphocytes, spleen and kidney. In addition, it is regulated by the tumor suppressor p53. TNFRSF10B is the receptor for the cytotoxic ligand TNFSF10/TRAIL. It

A DRUG SCREENING EXPERT

promotes the activation of NF-kappa-B and is essential for ER stress-induced apoptosis. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases mediating apoptosis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481