

Nectin-4 Protein, Human, Recombinant (aa 32-351, hFc)

General Information

| | |
|-----------------------|---|
| Synonyms: | PRR4;PVRL4;LNIR;Nectin-4;Ig superfamily receptor LNIR;Poliovirus receptor-related protein 4 |
| Protein Construction: | Gly32-Val351 |
| Species: | Human |
| Expression Host: | HEK293 Cells |
| Accession: | Q96NY8 |
| Molecular Weight: | 75-90 KDa (reducing condition) |
| AA Sequence: | Gly32-Val351 |

QC Testing

| | |
|----------------------|---|
| Biological Activity: | 1. Loaded Anti-Human Nectin-4 mAb-mFc on AMQ Biosensor, can bind Human Nectin-4-Fc with an affinity constant of 0.86 nM as determined in BLI assay. (Regularly tested) 2. Immobilized Anti-Human Nectin-4 mAb-mFc at 2µg/ml (100 µl/well) can bind Human Nectin-4-Fc . The ED50 of Human Nectin-4-Fc is 2.79 ng/ml. (Regularly tested) |
| Purity: | Greater than 95% as determined by reducing SDS-PAGE. Greater than 95% as determined by SEC-HPLC. |
| Endotoxin: | < 0.1 ng/µg (1 EU/µg) as determined by LAL test. |
| Formulation: | Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM PB, 150 mM NaCl, pH 7.4. |

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Nectin-4 (PVRL4) is a type I transmembrane glycoprotein which belongs to the nectin family of Ig superfamily proteins. It contains two Ig-like C2-type domains and one Ig-like V-type domain. PVRL4 seems to be involved in

cell adhesion through trans-homophilic and -heterophilic interactions, the latter including specifically interactions with nectin-1. It does not act as receptor for alpha-herpesvirus entry into cells. It is predominantly expressed in placenta, the embryo and breast carcinoma. But it is not detected in normal breast epithelium. The soluble form is produced by proteolytic cleavage at the cell surface (shedding), probably by ADAM17. Mutations in this gene are the cause of ectodermal dysplasia-syndactyly syndrome type 1, an autosomal recessive disorder.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481