Data Sheet (Cat.No.TMPJ-00330)

TargetMol

GPC1 Protein, Human, Recombinant (His)

General Information

Synonyms:	Glypican-1;GPC1
Protein Construction:	Asp24-Thr529
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P35052
Molecular Weight:	65-70 KDa (reducing condition)
AA Sequence:	Asp24-Thr529

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The Glypicans are a small multigene family of GPI-linked proteoglycans that play a key role in growth factor signaling. Human Glypican 1 (GPC1) is synthesized as a 558 amino acid (aa) preproprecursor that contains a 23 aa signal sequence, a 507 aa mature segment, and a 28 aa C-terminal prosegment. There are two potential N-linked and four potential O-linked sites for glycosylation or glycanation. There are potentially two heparan sulfate (HS) modifications on GPC1 that could contribute to a native molecular weight of approximately 200 kDa. Mature

human GPC1 shares 91% aa identity with mature mouse GPC1. Cells known to express GPC1 include neurons, smooth and skeletal muscle cells, keratinocytes, osteoblasts, Schwann cells, immature dendritic cells, and tumor, plus tumorassociated vascular endothelial cells. The function of GPC1 is complex and varied. As a proteoglycan, it appears to make use of its HS adduct to impact select growth factor activity. This is accomplished by having juxtramembrane HS attachment sites, and a flexible, GPI-linkage.

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481