

CD30/TNFRSF8 Protein, Human, Recombinant (aa 19-379, His)

General Information

Synonyms:	Lymphocyte activation antigen CD30;TNFRSF8;Tumor necrosis factor receptor superfamily member 8;Ki-1 antigen;CD30L receptor;CD30
Protein Construction:	Phe19-Lys379
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P28908
Molecular Weight:	60-95 KDa (reducing condition)
AA Sequence:	Phe19-Lys379

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM Tris-HCl, 150 mM NaCl, pH 7.5.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD30, also known as TNFRSF8, is a cell membrane protein of the tumor necrosis factor receptor family, which regulates proliferation/apoptosis and antibody responses. CD30 is expressed by activated, but not by resting, T and B cells. Aberrant expression of CD30 by mastocytosis mast cells and interaction with its ligand CD30L (CD153) appears to play an important role in the pathogenesis and clinical presentation of systemic mastocytosis. CD30

A DRUG SCREENING EXPERT

has been considered as a specific diagnostic biomarker of anaplastic large cell lymphoma (ALCL) and classical Hodgkin lymphoma (cHL). CD30 is also a biomarker used for targeted therapy by an antibody–drug conjugate.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481