

## IGF2BP-2 Protein, Human, Recombinant (His)

### General Information

Synonyms:	IGF-II mRNA-Binding Protein 2;IGF2 mRNA-Binding Protein 2;Hepatocellular Carcinoma Autoantigen p62;IGF2BP2;VICKZ2;IMP-2;VICKZ Family Member 2;IMP2;Insulin-Like Growth Factor 2 mRNA-Binding Protein 2
Protein Construction:	Met1-Thr220
Species:	Human
Expression Host:	E. coli
Accession:	Q9Y6M1
Molecular Weight:	43 KDa (reducing condition)
AA Sequence:	Met1-Thr220

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 90% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Insulin-Like Growth Factor 2 mRNA-Binding Protein 2 (IGFBP2) belongs to the RRM IMP/VICKZ family. IGFBP2 is a cytoplasmic protein and contains four KH domains and two RRM (RNA recognition motif) domains. IGF2BP2 binds to the 5'-UTR of the Insulin-Like Growth Factor 2 (IGF2) mRNA. This binding is isoform-specific. IGF2BP2 may regulate translation of target mRNAs. Genetic variation at the IGF2BP2 gene has been associated with type 2

diabetes (T2D) by genome-wide association studies and by replication analyses.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481