

GHR/Growth Hormone R Protein, Mouse, Recombinant (hFc)

General Information

Synonyms:	GHBP;GHR;Somatotropin receptor;growth hormone receptor;growth hormone binding protein;GH receptor;serum binding protein
Protein Construction:	Met1-Gln273
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q3UP14
Molecular Weight:	65-90 KDa (reducing condition)
AA Sequence:	Met1-Gln273

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Growth hormone receptor is a transmembrane receptor for growth hormone (GH). GH is a single-chain polypeptide that is mainly synthesized and released from the anterior pituitary gland and plays essential roles in growth, development and metabolism. GH exerts its physiological actions via GH binding to its receptor in its extracellular domain. Binding of growth hormone to the receptor leads to receptor dimerization and the activation of an intra- and intercellular signal transduction pathway leading to growth. Growth hormone receptor has been

shown to interact with SGTA, PTPN11, Janus kinase 2, Suppressor of cytokine signaling 1 and CISH.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481