

## FGFR2IIIb Protein, Human, Recombinant (aa 22-378, hFc)

### General Information

Synonyms:	FGF R2b;KGFR;BFR-1;FGFR2 $\beta$ ;FGFR2b;FGFR2;FGFR2 beta;CD332;FGFR2IIIb
Protein Construction:	Arg22-Glu378
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P21802-3
Molecular Weight:	85-120 KDa (reducing condition)
AA Sequence:	Arg22-Glu378

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/ $\mu$ g (1 EU/ $\mu$ g) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 $\mu$ m filter, containing PBS, pH 7.4.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100  $\mu$ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

FGFR2, also known as CD332, belongs to the fibroblast growth factor receptor subfamily where amino acid sequence is highly conserved between members and throughout evolution. FGFR2 acts as cell-surface receptor for fibroblast growth factors and plays an essential role in the regulation of cell proliferation, differentiation, migration and apoptosis, and in the regulation of embryonic development. Four distinct genes encoding closely related FGF receptors, FGF R1 - 4, are known. A frequent splicing event involving FGF R1 and 2 results in receptors

## A DRUG SCREENING EXPERT

---

containing all three Ig domains, referred to as the alpha isoform, or only IgII and IgIII, referred to as the beta isoform. Only the alpha isoform has been identified for FGF R3 and FGF R4. FGFR2 signaling is down-regulated by ubiquitination, internalization and degradation. Mutations that lead to constitutive kinase activation or impair normal FGFR2 maturation, internalization and degradation lead to aberrant signaling. Over-expressed FGFR2 promotes activation of STAT1.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481