

DLK1 Protein, Human, Recombinant (His)

General Information

Synonyms:	pG2;Protein Delta Homolog 1;Protein δ Homolog 1;DLK;DLK-1;DLK1
Protein Construction:	Ala24-Pro297(Ser260Asn)
Species:	Human
Expression Host:	HEK293 Cells
Accession:	AAH13197.1
Molecular Weight:	35-45 KDa (reducing condition)
AA Sequence:	Ala24-Pro297(Ser260Asn)

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/ μ g (1 EU/ μ g) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing 20 mM PB, 150 mM NaCl, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Protein Delta Homolog 1 (DLK-1) is a transmembrane protein which contains a signal peptide, an extracellular domain with six tandem epidermal growth factor (EGF)-like domains, a single pass transmembrane domain, and a short cytoplasmic tail. It is found within the stromal cells in close contact to the vascular structure of placental villi, yolk sac, fetal liver, adrenal cortex and pancreas and in the beta cells of the islets of Langerhans in the adult pancreas. In addition, it is detected in some forms of neuroendocrine lung tumor tissue. DLK-1 may have a

important role in neuroendocrine differentiation.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481