Data Sheet (Cat.No.TMPJ-00531)

TargetMol

TSLPR Protein, Mouse, Recombinant (His)

General Information

Synonyms:	Cytokine receptor-like 2;Thymic stromal-derived lymphopoietin receptor;CRLF2;cytokine receptor-like factor 2;TSLPR;Thymic stromal lymphopoietin protein receptor;P2RY8/CRLF2 fusion;CRL2;IL-XR;ILXR;CRL2 cytokine receptor;TSLP receptor
Protein Construction:	Ala20-Leu233
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	AAH23788.1
Molecular Weight:	35-40 KDa (reducing condition)
AA Sequence:	Ala20-Leu233

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 150 mM NaCl, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The cytokine thymic stromal lymphopoietin receptor (TSLPR) is consisting of a common γ receptor-like chain (TSLPR-γ) and a common interleukin 7 (IL-7) Rα chain that belongs to the type 1 cytokine receptor family. Transfection of TSLPR cDNA result in only low affinity binding, while cotransfection of the IL-7Rα chain cDNA shows

high affinity binding. TSLP and TSLPR play a critical role in the initiation of allergic diseases in mice. The TSLP R cDNA encodes a transmembrane receptor containing 370 amino acids (aa) with two potential N-linked glycosylation sites and a cytoplasmic domain of 104 aa including a single tyrosine residue. TSLPR can mediate signaling of the signal transducer and activator of transcription 5 (Stat5) by TSLP. TSLP R is broadly expressed in the immune and hematopoietic cells, particularly in hematopoietic progenitors and myeloid cells.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481