

hFcγR4 Protein, Mouse, Recombinant (His)

General Information

Synonyms:	Low Affinity Immunoglobulin γ Fc Region Receptor IV;FcγR4;Low Affinity Immunoglobulin Gamma Fc Region Receptor IV;CD16-2
Protein Construction:	Gly21-Gln203
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	AAH27310.1
Molecular Weight:	25-35 KDa (reducing condition)
AA Sequence:	Gly21-Gln203

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

FcγR4, also known as CD16-2, is one of the receptors for Fc region of IgG which involve in immune responses. FcγR4 mainly functions in cellular response to lipopolysaccharide, NK T cell proliferation, regulation of sensory perception of pain, wound healing etc. Three groups are included for Fc γ receptors (FcR), and they are Fc γ RI (CD64), Fc γ RII (CD32), and Fc γ RIII (CD16). Among these, CD64 possess high affinity even for monomeric IgG, while CD32 and CD16 display a relative lower affinity for IgG. Genes encodes these receptors are diverse differing

A DRUG SCREENING EXPERT

by species and cell types. The aggregation of FcR having immunoreceptor tyrosine-based activation motifs (ITAMs) activates sequentially src family tyrosine kinases and syk family tyrosine kinases that connect transduced signals to common activation pathways shared with other receptors. FcR with ITAMs elicit cell activation, endocytosis, and phagocytosis. Fc γ R3 belongs to Fc γ R3 (CD16) group.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481