

ANGPTL7 Protein, Human, Recombinant (His)

General Information

Synonyms:	CDT6;Cornea-derived transcript 6 protein;Angiopoietin-like protein 7;Angiopoietin-like factor;ANGPTL7;Angiopoietin-related protein 7
Protein Construction:	Gln27-Pro346
Species:	Human
Expression Host:	HEK293 Cells
Accession:	O43827
Molecular Weight:	34745-55 KDa (reducing condition)
AA Sequence:	Gln27-Pro346

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 6% Trehalose, 4% Mannitol, 100 mM NaCl, 0.05% Tween 80, pH6.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Angiopoietin-like 7 (ANGPTL7) is a secreted glycoprotein that is structurally related to the angiopoietins. Members of this protein family contain an N-terminal coiled coil domain and a C-terminal fibrinogen-like domain. ANGPTL7 shares 89% aa sequence identity with mouse and rat ANGPTL7. It is secreted as a 45-50kDa monomer that forms disulfide-linked homotrimers and tetramers via the coiled coil domain. ANGPTL7 is expressed in the corneal

stroma, trabecular meshwork, and sclera and is elevated in glaucoma aqueous humor. Its production is up-regulated in trabecular meshwork cells by glucocorticoids and TGF- β and in cartilage by TNF- α . Overexpression of ANGPTL7 in trabecular meshwork cells inhibits the production of collagen and proteoglycans. When overexpressed in tumor cells it promotes collagen and proteoglycan deposition but inhibits tumor xenograft progression and tumor angiogenesis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481