

## Coagulation factor X/F10 Protein, Mouse, Recombinant (His)

### General Information

Synonyms:	Stuart factor;Coagulation factor X;F10
Protein Construction:	Gly21-Asn481
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	O88947
Molecular Weight:	50-60&20-28 KDa (reducing condition)
AA Sequence:	Gly21-Asn481

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM MES, 150 mM NaCl, 1 mM CaCl <sub>2</sub> , pH 7.5.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Mouse coagulation factor X / F10 a member of the peptidase S1 family. The mature F10 is composed mostly of two EGF-like domains, one Gla gamma-carboxy-glutamate domain and one peptidase S1 domain. Factor Xa is a vitamin K-dependent plasma protease that converts prothrombin to thrombin in the presence of factor Va, calcium and phospholipid during blood clotting. The two chains of F10 are formed from a single-chain precursor by the excision of two Arg residues. A single-chain precursor is initially synthesized in the liver. The light and heavy

chains are linked together by disulfide bonds. The light chain contains a Gla and two EGF-like domains. The heavy chain corresponds to the serine protease domain. It can form a heterodimer with SERPINA5.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481