

## GADD45G Protein, Human, Recombinant (His)

### General Information

Synonyms:	Cytokine-Responsive Protein CR6;Growth Arrest and DNA Damage-Inducible Protein GADD45 Gamma;Growth Arrest and DNA Damage-Inducible Protein GADD45 $\gamma$ ;DDIT-2;DNA Damage-Inducible Transcript 2 Protein;CR6;DDIT2;GADD45G
Protein Construction:	Met1-Glu159
Species:	Human
Expression Host:	E. coli
Accession:	O95257
Molecular Weight:	21 KDa (reducing condition)
AA Sequence:	Met1-Glu159

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/ $\mu$ g (1 EU/ $\mu$ g) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 $\mu$ m filter, containing 20 mM Tris-HCl, 150 mM NaCl, pH 8.0.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100  $\mu$ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Growth Arrest and DNA Damage-Inducible Protein GADD45 Y (GADD45G) is a nuclear protein which belongs to the GADD45 family. GADD45G is highly expressed in placenta. GADD45G interacts with various proteins whose transcript levels are increased following stressful growth arrest conditions and treatment with DNA-damaging

## A DRUG SCREENING EXPERT

---

agents. GADD45G responds to environmental stresses by mediating activation of the p38/JNK pathway via MTK1/MEKK4 kinase. GADD45G is also involved in the regulation of growth and apoptosis. GADD45G inhibits cell growth and differentiation by androgens. The mRNA expression is down-regulated in hepatocellular carcinoma.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481