

EGR1 Protein, Human, Recombinant (His)

General Information

Synonyms:	zinc finger protein 225;NGFI-A;Zif268;Early growth response protein 1;EGR-1;nerve growth factor-induced protein A
Protein Construction:	Gln282-Ser433
Species:	Human
Expression Host:	E. coli
Accession:	P18146
Molecular Weight:	19 KDa (reducing condition)
AA Sequence:	Gln282-Ser433

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 150 mM NaCl, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

EGR-1 belongs to the EGR family of C2H2-type zinc finger proteins. It is a nuclear protein and functions as a transcriptional regulator. EGR-1 recognizes and binds to the DNA sequence 5'-CGCCCCGC-3'(EGR-site).The products of target genes it activates are required for differentiation and mitogenesis. Studies suggest this is a tumor suppressor gene. EGR-1 has a distinct pattern of expression in the brain, and its induction has been shown

A DRUG SCREENING EXPERT

to be associated with neuronal activity. Several studies suggest it has a role in neuronal plasticity. EGR-1 has also been found to regulate the expression of synaptobrevin II (a protein important for synaptic exocytosis).

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481