

## NFYA Protein, Human, Recombinant (GST)

### General Information

Synonyms:	Nuclear Transcription Factor Y Subunit Alpha; Nuclear Transcription Factor Y Subunit A; NFYA; CAAT Box DNA-Binding Protein Subunit A; Nuclear Transcription Factor Y Subunit $\alpha$ ; NF-YA
Protein Construction:	Met1-Ser318
Species:	Human
Expression Host:	E. coli
Accession:	P23511-2
Molecular Weight:	55 KDa (reducing condition)
AA Sequence:	Met1-Ser318

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/ $\mu$ g (1 EU/ $\mu$ g) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 $\mu$ m filter, containing 20 mM PB, 150 mM NaCl, pH 7.2.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100  $\mu$ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Nuclear Transcription Factor Y Subunit  $\alpha$  (NFYA) is a member of the NFYA/HAP2 subunit family. NFYA functions as a heterotrimeric transcription factor, which is composed of three components, NF-YA, NF-YB and NF-YC, binds to CCAAT motifs in the promoter regions in a variety of genes. NFYA forms a highly conserved transcription factor which stimulates the transcription of various genes by recognizing and binding to a CCAAT motif in promoters, for

example in type 1 collagen, albumin and beta-actin genes.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481