

PEA15 Protein, Human, Recombinant

General Information

Synonyms:	15 kDa Phosphoprotein Enriched in Astrocytes;PEA15;PED;Astrocytic Phosphoprotein PEA-15; Phosphoprotein Enriched in Diabetes
Protein Construction:	Met1-Ala130
Species:	Human
Expression Host:	E. coli
Accession:	Q15121
Molecular Weight:	12–16 KDa (reducing condition)
AA Sequence:	Met1-Ala130

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 150 mM NaCl, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Astrocytic phosphoprotein PEA-15 (PEA15) is a death effector domain (DED)-containing protein. PEA15 is mainly expressed in the central nervous system, principally in astrocytes. Increased PEA15 levels affect tumorigenesis and cancer progression. PEA15 is overexpressed in breast cancers and gliomas as well as in type 2 diabetes. PEA15 blocks Ras-mediated inhibition of integrin activation and modulates the ERK MAP kinase cascade. PEA15 also

A DRUG SCREENING EXPERT

inhibits RPS6KA3 activities by holding it in the cytoplasm. In addition, PEA15 inhibits both TNFRSF6 and TNFRSF1A mediated CASP8 activity and apoptosis. At present, PEA15 expression is also a significant prognostic marker in ovarian cancer.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481