

ABCB5 Protein, Human, Recombinant (Trx)

General Information

Synonyms:	ABCB5 P-gp;ATP-binding cassette sub-family B member 5;ABCB5;P-glycoprotein ABCB5
Protein Construction:	Ile141-Val247
Species:	Human
Expression Host:	E. coli
Accession:	Q2M3G0
Molecular Weight:	30 KDa (reducing condition)
AA Sequence:	Ile141-Val247

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM PB, 150 mM NaCl, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

ATP-binding cassette sub-family B member 5(ABCB5) is a plasma membrane-spanning protein. ABCB5 is principally expressed in physiological skin and human malignant melanoma. ABCB5 has been suggested to regulate skin progenitor cell fusion and mediate chemotherapeutic drug resistance in stem-like tumor cell subpopulations in human malignant melanoma. It is commonly over-expressed on circulating melanoma tumour cells. Furthermore, the ABCB5+ melanoma- initiating cells were demonstrated to express FLT1 (VEGFR1) receptor

tyrosine kinase which was functionally required for efficient xenograft tumor formation, as demonstrated by shRNA knockdown experiments.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481