

## TGFBR1 Protein, Mouse, Recombinant (hFc)

## General Information

Synonyms:	ALK-5;ALK-5ALK5;TGFβRI;tβR-I;LDS1A;TGFbetaRI;TGFB1R1;activin A receptor type II-like kinase, 53kD;TGF-β receptor type I;LDS2A;TGF-bRI;ACVRLK4;TGFR-1;SKR4;TGFBR1;TGF-beta receptor type I;AAT5;tbetaR-I
Protein Construction:	Leu30-Glu125
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q64729
Molecular Weight:	40-60 KDa (reducing condition)
AA Sequence:	Leu30-Glu125

## QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

## Preparation and Storage

## Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

## Stability &amp; Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

## Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

## Protein Background

TGF-beta RI, also called ALK-5, is an approximately 55 kDa type I transmembrane serine/threonine receptor kinase. In the presence of TGF-beta, TGF-beta RI forms a complex with, and is phosphorylated by, TGF-beta RII. Phosphorylated TGF-beta RI can then transiently bind and phosphorylate Smad2 and Smad3. TGF-beta functions as a tumor suppressor by inhibiting the cell cycle in the G1 phase. Administration of TGF-beta is able to protect

## A DRUG SCREENING EXPERT

---

against mammary tumor development in transgenic mouse models in vivo. Disruption of the TGF-beta/SMAD pathway has been implicated in a variety of human cancers, with the majority of colon and gastric cancers being caused by an inactivating mutation of TGF-beta RII. TGF-beta RI is likely important during development, since mice deficient for TGF-beta RI die at midgestation with severe defects in vascular development of the yolk sac and placenta, and an absence of circulating red blood cells. Furthermore, TGF-beta RI appears to be involved in proper lymphatic network development.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481