

EDA2R Protein, Mouse, Recombinant (His)

General Information

Synonyms:	EDA-A2R;Tumor necrosis factor receptor superfamily member XEDAR;EDA-A2 receptor; XEDAR;EDAR2;X-linked ectodysplasin-A2 receptor;TNFRSF27;Ectodysplasin A2 receptor; Tumor necrosis factor receptor superfamily member 27;EDAA2R
Protein Construction:	Met1-Thr138
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	Q8BX35
Molecular Weight:	26 KDa (reducing condition)
AA Sequence:	Met1-Thr138

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:
Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Tumor necrosis factor receptor superfamily member 27, also known as XEDAR and EDA2R, is a type III transmembrane protein of the TNFR superfamily. EDA2R consists of extracellular domain (ECD) with 3 cysteine-rich repeats and a single transmembrane domain but lacks an N-terminal signal peptide. EDA2R is widely expressed, notably in embryonic basal epidermal cells and maturing hair follicles. Even though it does not contain a

cytoplasmic death domain, EDA2R can associate with Fas and induce EDA-A2 dependent apoptosis. Its transcription is directly induced by p53, and its mediated cell death is p53 dependent. It is down-regulated in breast, colon, and lung cancers, particularly in cases with p53 mutations. It also plays a role in EDA-A2 induced skeletal muscle degeneration and osteoblast differentiation. Mutations in the EDA gene are associated with the X-linked form of Hypohidrotic Ectodermal Dysplasia (HED), a disease typically characterized by abnormal hair, teeth and sweat glands.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481