

CD99L2 Protein, Human, Recombinant (hFc)

General Information

Synonyms:	CD99 Antigen-Like Protein 2;CD99;CD99L2;MIC2L1;MIC2-Like Protein 1
Protein Construction:	Asp26-Ala188
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q8TCZ2
Molecular Weight:	60-90 KDa (reducing condition)
AA Sequence:	Asp26-Ala188

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS, pH 7.4.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD99 Antigen-Like Protein 2 (CD99L2) belongs to the CD99 family. CD99L2 is a single-pass type I membrane protein and expressed in many tissues, with low expression in thymus. CD99L2 plays a role in a late step of leukocyte extravasation helping cells to overcome the endothelial basement membrane. CD99L2 and CD99 are involved in trans-endothelial migration of neutrophils in vitro and in the recruitment of neutrophils into inflamed peritoneum. A similar protein in mouse functions as an adhesion molecule during leukocyte extravasation.

Alternate splicing results in multiple transcript variants.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481