

Caspase-10 Protein, Human, Recombinant (His)

General Information

Synonyms:	CASP10;FAS-Associated Death Domain Protein Interleukin-1B-Converting Enzyme 2;CASP-10;ICE-Like Apoptotic Protease 4;Caspase-10;MCH4;Apoptotic Protease Mch-4;FLICE2
Protein Construction:	Val220-Ile480
Species:	Human
Expression Host:	E. coli
Accession:	Q92851-4
Molecular Weight:	33 KDa (reducing condition)
AA Sequence:	Val220-Ile480

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 70% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Supplied as a 0.2 μm filtered solution of 20 mM Tris-HCl, 8% Sucrose, 1 mM DTT, 0.05% Tween 80, pH8.5.

Preparation and Storage

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

Proteins are shipped with blue ice.

Protein Background

Caspase-10 (CASP10) is a 521 amino acid protein member of the Cysteine-Aspartic Acid Protease (Caspase) family. CASP10 contains two DED (Death Effector) domains and is detectable in most tissues. CASP10 cleavage by Granzyme B and autocatalytic activity generate the two active subunits: Caspase-10 subunit p23/17, Caspase-10 subunit p12. Caspases are a family of cytosolic aspartate-specific cysteine proteases involved in the execution-phase of cell apoptosis, the initiation and execution. Human caspases can be subdivided into three functional groups: cytokine activation (caspase-1, -4, -5, and -13), apoptosis initiation (caspase-2, -8, -9, -and -10), and apoptosis execution (caspase-3, -6, and -7). CASP10 cleaves and activates caspases 3 and 7, but itself is processed by caspase 8. Defects in CASP10 are associated with apoptosis defects seen in type II autoimmune lymphoproliferative syndrome.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481