

## CLIC5 Protein, Human, Recombinant (His)

### General Information

Synonyms:	Chloride Intracellular Channel Protein 5;CLIC5
Protein Construction:	Met1-Ser251
Species:	Human
Expression Host:	E. coli
Accession:	Q9NZA1
Molecular Weight:	32 KDa (reducing condition)
AA Sequence:	Met1-Ser251

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM Tris-HCl, 150 mM NaCl, 0.01% Tween 80, 0.01% TritonX-100, pH8.0.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Chloride Intracellular Channel Protein 5 (CLIC5) is a single-pass membrane protein which belongs to the chloride channel CLIC family. It contains one GST C-terminal domain. Chloride intracellular channels are involved in chloride ion transport within various subcellular compartments. CLIC5 can insert into membranes and form selective ion channels regulated by actin that may transport chloride ions. It may play a role in the regulation of transepithelial ion absorption and secretion. CLIC5 specifically associates with the cytoskeleton of placenta

microvilli. CLIC5 is required for the development and/or maintenance of the proper glomerular endothelial cell and podocyte architecture.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481