

## ASF1A Protein, Human, Recombinant (His, T7)

### General Information

Synonyms:	CIA;Anti-Silencing Function Protein 1 Homolog A;ASF1A;Histone Chaperone ASF1A;hAsf1;hAsf1a;CCG1-Interacting Factor A;hCIA
Protein Construction:	Met1-Met204
Species:	Human
Expression Host:	E. coli
Accession:	Q9Y294
Molecular Weight:	22 KDa (reducing condition)
AA Sequence:	Met1-Met204

### QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 95% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 0.1 ng/μg (1 EU/μg) as determined by LAL test.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing 20 mM Tris-HCl, 1 mM DTT, 150 mM NaCl, pH 8.0.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months.

Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

Human Histone Chaperone ASF1A (ASF1A) belongs to the H3/H4 family of histone chaperone proteins. ASF1A is ubiquitously expressed in many cells and tissues, interacting with histones H3 and H4. ASF1A cooperates with Chromatin Assembly Factor 1 to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly. In addition, ASF1A is necessary for the formation of senescence-

associated heterochromatin foci (SAHF) and efficient senescence-associated cell cycle exit.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481