

MKI67 Protein, Human, Recombinant (GST)

General Information

Synonyms:	Ki-67;Antigen Ki67;MKI67;KIA;antigen KI-67;MIB-1;MIB-;Ki67;PPP1R105
Protein Construction:	Met1-Pro120
Species:	Human
Expression Host:	E. coli
Accession:	P46013
Molecular Weight:	38 KDa (reducing condition)
AA Sequence:	Met1-Pro120

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	Greater than 80% as determined by reducing SDS-PAGE. (QC verified)
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM Tris-HCl, 8% Sucrose, 0.05% Tween 80, pH 8.0.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

MKI67 also known as Ki67, is a 350-400 kDa nuclear protein that belongs to a molecular group comprised of mitotic chromosome-associated proteins. MKI67 contains 1 FHA domain and plays a key role in cell proliferation. MKI67 is contextually expressed, being potentially found in all cells that are not in the Go phase of the cell cycle. Thus, MKI67 qualifies as a cell proliferation marker. It is also associated with ribosomal RNA transcription. Inactivation of antigen MKI67 leads to inhibition of ribosomal RNA synthesis.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481