

CD94 Protein, Human, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	NK cell receptor;CD94;KP43;KLRD1
Protein Construction:	Ser34-Ile179
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q13241-1
Molecular Weight:	19.8 kDa (predicted). Due to glycosylation, the protein migrates to 35-45 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

CD94 is an approximately 25 kDa type 2 transmembrane protein that plays an important role in regulating natural killer (NK) cell activation. CD94 plays a role as a receptor for the recognition of MHC class I HLA-E molecules by NK cells and some cytotoxic T-cells.

Reference

Walsh CE, et al. Differential Expression of NK Receptors CD94 and NKG2A by T Cells in Rheumatoid Arthritis Patients in Remission Compared to Active Disease[J]. PLOS ONE, 2011, 6.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481