

CD32B/Fcgr2b Protein, Human, Recombinant (His & Avi)

General Information

Synonyms:	CDw32;Fc γ RIIB;Fc gamma RIIB;Fc γ RII;FCG2;IGFR2;Fc gamma RII;FcGR;CD32b;FCGR21;FCGR2;CD32;FCGR2A;CD32A
Protein Construction:	Ala46-Pro217
Species:	Human
Expression Host:	HEK293 Cells
Accession:	P31994-1
Molecular Weight:	22.5 kDa (predicted). Due to glycosylation, the protein migrates to 32-37 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Rituximab captured on CM5 Chip via Protein A can bind Human Fc gamma RIIB, His Tag with an affinity constant of 3.46 μ M as determined in SPR assay.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The Fc gamma Rs have been divided into three classes based on close relationships in their extracellular domains; these groups are designated Fc gamma RI (also known as CD64), Fc gamma RII (CD32), and Fc gamma RIII (CD16). Each group may be encoded by multiple genes and exist in different isoforms depending on species and cell type.

Reference

Engelhardt W, et al. DISTRIBUTION, INDUCIBILITY AND BIOLOGICAL FUNCTION OF THE CLONED AND EXPRESSED HUMAN BETA-FC RECEPTOR-II[J]. European Journal of Immunology, 2010, 20(6):1367-1377.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481