

uPAR/PLAUR Protein, Mouse, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	U-PAR;PLAUR;MO3;CD87;uPAR
Protein Construction:	Leu24-Gly298
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P35456-1
Molecular Weight:	32.92 kDa (predicted). Due to glycosylation, the protein migrates to 55-65 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Biotinylated Mouse uPAR, His Tag immobilized on CM5 Chip can bind Human PLAUR, His Tag with an affinity constant of 6.35 nM as determined in SPR assay.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The receptor (u-PAR) for urokinase plasminogen activator (u-PA) is a three-domain protein, GPI-anchored to the cell surface, which focuses the enzymatic activity of u-PA, and allows the cell surface activation of plasminogen. Regulation of the activity of u-PA is also mediated by u-PAR.

Reference

Blasi F. The urokinase receptor and cell migration. Semin Thromb Hemost. 1996;22(6):513-6. doi: 10.1055/s-2007-999053. PMID: 9122717.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481