

LILRA2/CD85h/ILT1 Protein, Human, Recombinant (His & Avi), Biotinylated

General Information

Synonyms:	LIR7;ILT-1;ILT1;LILRA2;CD85h;ILT1CD85H
Protein Construction:	Gly24-Asn449
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q8N149-1
Molecular Weight:	49.9 kDa (predicted). Due to glycosylation, the protein migrates to 70-80 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

LILRA2, also known as ILT1, CD85h, and LIR7, is an approximately 70 kDa variably glycosylated transmembrane protein that regulates immune cell activation. Mature human LILRA2 consists of a 426 amino acid (aa) extracellular domain (ECD) with 4 Ig-like domains, a 21 aa transmembrane segment, and a 13 aa cytoplasmic tail. LILRA2 is part of the innate immune responses against microbial infection.

Reference

Lu H K, et al. LILRA2 Selectively Modulates LPS-Mediated Cytokine Production and Inhibits Phagocytosis by Monocytes[J]. PLoS ONE, 2012, 7(3):e33478-.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481