

IFNAR1 Protein, Mouse, Recombinant (His & Avi)

General Information

Synonyms:	IFN- α -REC;IFN-R-1;IFNAR1;IFN-alpha/ β R1;IFRC;IFN α / β R1;IFN- α / β R1;IFNAR;IFNBR;AVP;IFN-alpha-REC;IFNbR1;CRF2-1;IFN-aR1;IFN-alpha/beta R1
Protein Construction:	Glu27-Thr429
Species:	Mouse
Expression Host:	HEK293 Cells
Accession:	P33896
Molecular Weight:	46.9 kDa (predicted). Due to glycosylation, the protein migrates to 67-70 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Mouse IFNAR1, His Tag captured on CM5 Chip via anti-his antibody can bind Human IFN alpha 1, hFc Tag with an affinity constant of 3.61 nM as determined in SPR assay.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

IFN-alpha / beta R1, also known as IFNAR1, belongs to the class II cytokine receptor family of proteins. Class II cytokine receptors form heterodimeric receptor complexes that mediate class II cytokine signals. Subunits of the different receptor complexes are shared and serve multiple functions. Functions in general as heterodimer with IFNAR2. Type I interferon binding activates the JAK-STAT signaling cascade, and triggers tyrosine phosphorylation

of a number of proteins including JAKs, TYK2, STAT proteins and the IFNR alpha- and beta-subunits themselves.

Reference

Teunissen P F A, et al. MAb therapy against the IFN- α/β receptor subunit 1 stimulates arteriogenesis in a murine hindlimb ischaemia model without enhancing atherosclerotic burden[J]. Cardiovascular Research, 2015, 107(2): 255-266.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481