

AXL Protein, Cynomolgus, Recombinant (His)

General Information

Synonyms:	Tyro7;JTK11;AI323647;AXL oncogene;EC 2.7.10.1;EC 2.7.10;UFO;ARK;Axl
Protein Construction:	Glu33-Pro449
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A1D5Q330
Molecular Weight:	46.3 kDa (predicted). Due to glycosylation, the protein migrates to 68-70 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	<ol style="list-style-type: none">1. Immobilized Cynomolgus AXL, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-AXL Antibody, hFc Tag with the EC50 of 8.8ng/ml determined by ELISA (QC Test).2. Cynomolgus Axl, His Tag immobilized on CM5 Chip can bind Cynomolgus GAS6, His Tag with an affinity constant of 1.70 nM as determined in SPR assay.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Axl, a member of the TAM (Tyro3, Axl, Mer) family, and its inhibitors can specifically break the kinase signaling nodes, allowing advanced patients to regain drug sensitivity with improved therapeutic efficacy. Overexpression

and activation of Axl receptor tyrosine kinase have been widely accepted to promote cell proliferation, chemotherapy resistance, invasion, and metastasis in several human cancers, such as lung, breast, and pancreatic cancers.

Reference

Zhu C, et al. AXL receptor tyrosine kinase as a promising anti-cancer approach: functions, molecular mechanisms and clinical applications. Mol Cancer. 2019 Nov 4;18(1):153. doi: 10.1186/s12943-019-1090-3. PMID: 31684958; PMCID: PMC6827209.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel: 781-999-4286 E_mail: info@targetmol.com Address: 34 Washington Street, Wellesley Hills, MA 02481