

BDCA-2 Protein, Cynomolgus, Recombinant (His)

General Information

Synonyms:	CD303;HECL;CLEC4C;DLEC;CLECSF7;CLECSF11;BDCA-2;BDCA2
Protein Construction:	Tyr48-Ile212
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A2K5UWP4
Molecular Weight:	20.55 kDa (predicted). Due to glycosylation, the protein migrates to 28-38 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Cynomolgus BDCA-2, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-BDCA-2 Antibody, hFc Tag with the EC50 of 4.8ng/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS, 300 mM L-arginine (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

BDCA-2, BDCA-3, and BDCA-4. In blood, BDCA-2 and BDCA-4 are expressed on CD11c(-) CD123(bright) plasmacytoid dendritic cells, whereas BDCA-3 is expressed on small population of CD11c() CD123(-) dendritic cells. All three Ags are not detectable on a third blood dendritic cell population, which is CD1c() CD11c(bright) CD123

(dim), or on any other cells in blood. BDCA-2 is completely down-regulated on plasmacytoid CD11c(-) CD123 (bright) dendritic cells, BDCA-2 is rapidly internalized at 37 degrees C after mAb labeling.

Reference

Dzionic A, et al. BDCA-2, BDCA-3, and BDCA-4: three markers for distinct subsets of dendritic cells in human peripheral blood. J Immunol. 2000 Dec 1;165(11):6037-46. doi: 10.4049/jimmunol.165.11.6037. PMID: 11086035.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481