

KLKB1 Protein, Cynomolgus, Recombinant (His)

General Information

Synonyms:	KLK3;Plasma kallikrein;KLKB1;PKK;Kininogenin;Fletcher factor
Protein Construction:	Gly20-Ala638
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A2K5VTJ9
Molecular Weight:	70.19 kDa (predicted). Due to glycosylation, the protein migrates to 70-90 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM NaAc, 150 mM NaCl (pH 5.0). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in 20mM NaAc, 150mM NaCl (pH 5.0). The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Plasma kallikrein, also known as Fletcher factor or kallikrein B1 (KLKB1), is a serine endopeptidase, like its homologs tissue kallikrein and kallikrein-related peptidases (KLKs). Its physiological role is to catalyze the release of kinins and other vasoactive peptides.

Reference

Abrera-Abeleda MA, et al. Mesangial immune complex glomerulonephritis due to complement factor D deficiency. *Kidney Int.* 2007 Jun;71(11):1142-7. doi: 10.1038/sj.ki.5002235. Epub 2007 Apr 4. PMID: 17410102.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481