

## CD46 Protein, Cynomolgus, Recombinant (His)

### General Information

Synonyms:	CD46;MCP;TLX;MIC10;AHUS2;TRA2.10
Protein Construction:	Cys35-Asp329
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	A0A2K5WCS2
Molecular Weight:	34.05 kDa (predicted). Due to glycosylation, the protein migrates to 50-70 kDa based on Tris-Bis PAGE result.

### QC Testing

Biological Activity:	Immobilized Cynomolgus CD46, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Anti-CD46 Antibody, hFc Tag with the EC50 of 0.67µg/ml determined by ELISA.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

### Preparation and Storage

#### Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

#### Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

#### Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

### Protein Background

CD46 was discovered in 1986 during a search for novel C3b-binding proteins. CD46 is expressed ubiquitously and functions as a co-factor in the factor I-mediated proteolytic cleavage of C3b and C4b. Its vital role in preventing complement deposition on host tissue is underpinned by the fact that deficiency of CD46 is a predisposing factor for numerous disease conditions arising from complement-mediated 'self-attack'.

Reference

Cardone J, et al. CD46 in innate and adaptive immunity: an update. Clin Exp Immunol. 2011 Jun;164(3):301-1doi: 10.1111/j.1365-2249.2011.04400.x. Epub 2011 Apr 13. PMID: 21488871; PMCID: PMC3087925.

**Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins**

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E\_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481