

TRAIL R1/DR4/TNFRSF10A Protein, Human, Recombinant (His & Avi)

General Information

Synonyms:	DR4;MGC9365;TNFRSF10A;TRAIL-R;TRAIL R1;CD261;APO2
Protein Construction:	Pro34-Asn239
Species:	Human
Expression Host:	HEK293 Cells
Accession:	O00220
Molecular Weight:	24.7 kDa (predicted). Due to glycosylation, the protein migrates to 25-30 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Human TRAIL R1, His Tag captured on CM5 Chip via anti-his antibody can bind Human TRAIL, No Tag with an affinity constant of 1.02 nM as determined in SPR assay.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/ μ g of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μ m filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μ g/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Tumor necrosis factor receptor superfamily member 10A (TNFRSF10A) is also known as TNF-related apoptosis-inducing ligand receptor 1 (TRAIL-R1), Death receptor 4 (DR4), CD261 and APO2, which belongs to TNF superfamily. The expression of apoptosis-inducing TRAIL-R1 and TRAIL-R2 and of the decoy receptors TRAIL-R3 and TRAIL-R4 was systematically studied in all developmental stages of peripheral B cells isolated from the blood and secondary lymphoid organs. Expression of TRAIL-Rs is modulated along development, with highest levels

observed in germinal center B cells.

Reference

Staniek J, et al. TRAIL-R1 and TRAIL-R2 Mediate TRAIL-Dependent Apoptosis in Activated Primary Human B Lymphocytes. Front Immunol. 2019 Apr 30;10:95 doi: 10.3389/fimmu.2019.0095 PMID: 31114586; PMCID: PMC6503035.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481