

APLP2 Protein, Human, Recombinant (His)

General Information

Synonyms:	APPL2;YWK-II;APPH;CDEI box-binding;CDEBP
Protein Construction:	Gly32-Ser692
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q06481-1
Molecular Weight:	77.16 kDa (predicted). Due to glycosylation, the protein migrates to 85-100 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Amyloid β precursor-like protein 2 (APLP2) has been determined to serve an important role in the progression of a number of cancer types. APLP2 expression was significantly associated with disease-specific survival (P<0.001). APLP2 may be used to potentially predict patient prognosis, and to guide clinical diagnosis and treatment in CCRCC.

Reference

Gao L, et al. Role of APLP2 in the prognosis and clinicopathology of renal cell carcinoma. *Oncol Lett.* 2019 Jan;17 (1):508-513. doi: 10.3892/ol.2018.9577. Epub 2018 Oct 15. PMID: 30655794; PMCID: PMC6313182.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481