

Cadherin 17/CDH17 Domain 6-7 Protein, Human, Recombinant (hFc)

General Information

Synonyms:	HPT-1;P130;MGC138218;LI-cadherin;FLJ26931;cadherin-16;Cdh17;CDH16;BILL-cadherin; Cadherin-17;MGC142024
Protein Construction:	Ser567-Gly777
Species:	Human
Expression Host:	HEK293 Cells
Accession:	Q12864
Molecular Weight:	49.29 kDa (predicted). Due to glycosylation, the protein migrates to 55-65 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Immobilized Human CDH17 Domain 6-7, His Tag at 5 µg/mL (100 µL/well) on the plate. Dose response curve for Biotinylated Anti-CDH17 Domain 6-7 Antibody, hFc Avi Tag with the EC50 of 0.19 µg/mL determined by ELISA.(QC Test)
Purity:	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC
Endotoxin:	< 1.0 EU/µg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 µm filter, containing 20 mM Tris, 150 mM NaCl (pH 8.0). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 µg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

Liver-intestine cadherin (CDH17) has been known to function as a tumor stimulator and diagnostic marker for almost two decades. In vivo studies showed CDH17 knockout resulted in apoptotic PC tumor death through

activating caspase-3 activity. Taken together, CDH17 functions as an oncogenic molecule critical to PC growth by regulating tumor apoptosis signaling pathways and CDH17 could be targeted to develop an anti-PC therapeutic approach.

Reference

Liu X, et al. Disruption of oncogenic liver-intestine cadherin (CDH17) drives apoptotic pancreatic cancer death. *Cancer Lett.* 2019 Jul 10;454:204-214. doi: 10.1016/j.canlet.2019.04.022. Epub 2019 Apr 17. PMID: 31004701.

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only · Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481