

IL-2 R gamma/CD132 Protein, Cynomolgus, Recombinant (His)

General Information

Synonyms:	SCIDX1;IL2RG;gammaC;γC;CIDX;IL-2 R γ/CD132;IL-2 R gamma;SCIDX;CD132;P64;IL-2 R γ;IMD4
Protein Construction:	Leu23-Asn254
Species:	Cynomolgus
Expression Host:	HEK293 Cells
Accession:	Q38JL2
Molecular Weight:	28.2 kDa (predicted). Due to glycosylation, the protein migrates to 50-68 kDa based on Tris-Bis PAGE result.

QC Testing

Biological Activity:	Activity has not been tested. It is theoretically active, but we cannot guarantee it. If you require protein activity, we recommend choosing the eukaryotic expression version first.
Purity:	> 95% as determined by Tris-Bis PAGE
Endotoxin:	< 1.0 EU/μg of the protein as determined by the LAL method.
Formulation:	Lyophilized from a solution filtered through a 0.22 μm filter, containing PBS (pH 7.4). Typically, 8% trehalose is incorporated as a protective agent before lyophilization.

Preparation and Storage

Reconstitution:

Reconstitute the lyophilized protein in distilled water. The product concentration should not be less than 100 μg/ml. Before opening, centrifuge the tube to collect powder at the bottom. After adding the reconstitution buffer, avoid vortexing or pipetting for mixing.

Stability & Storage:

It is recommended to store recombinant proteins at -20°C to -80°C for future use. Lyophilized powders can be stably stored for over 12 months, while liquid products can be stored for 6-12 months at -80°C. For reconstituted protein solutions, the solution can be stored at -20°C to -80°C for at least 3 months. Please avoid multiple freeze-thaw cycles and store products in aliquots.

Actual storage temperature shall be subject to the COA.

Shipping:

In general, lyophilized powders are shipped with blue ice, while solutions are shipped with dry ice.

Protein Background

The gamma chain of the high affinity functional human IL-2 receptor complex belongs to the hematopoietin receptor family. IL-2 R gamma is a 369 amino acid residue protein consisting of a 22 residue signal sequence, a 232 residue extracellular domain, a 29 residue transmembrane domain and an 86 residue cytoplasmic domain. Although IL-2 R gamma by itself does not bind IL-2 with any appreciable affinity, it is required for IL-2 receptor

signaling.

Reference

Smith K A. The structure of IL2 bound to the three chains of the IL2 receptor and how signaling occurs[J]. Medical Immunology, 2006, 5(1).

Inhibitor · Natural Compounds · Compound Libraries · Recombinant Proteins

This product is for Research Use Only· Not for Human or Veterinary or Therapeutic Use

Tel:781-999-4286 E_mail:info@targetmol.com Address:34 Washington Street,Wellesley Hills,MA 02481